

Well Child Care 18 Months

Nutrition

Family meals are important for your baby. Let him eat with you. This helps him learn. Don't . make mealtime a battle. Let your baby feed himself. Your child should use a spoon and drink from a cup now.

Development and Discipline

Children at this age should be learning many new words. You can help your child's vocabulary grow by showing and naming lots of things. Children have many different feelings and behaviors such as pleasure, anger, joy, curiosity, warmth, and assertiveness. It is important at this age to praise your child for doing things that you like. Make an effort to catch your children being good.

Toddlers often seem out of control, or overly stubborn or demanding. At this age, children often say "no" or refuse to do what you want them to do. Here are some good methods for helping children learn about rules and to keep them safe:

1. Child-proof the home. Go through every room in your house and remove anything that is either valuable, dangerous, or messy. Preventive child-proofing will stop many possible discipline problems. Don't expect a child not to get into things just because you say no.
2. Divert and substitute: If a child is playing with something you don't want him to have, replace it with another object or toy that he enjoys. This approach avoids a fight and does not place children in a situation where they will say "no."
3. Teach and lead. Have as few rules as necessary and enforce them. These rules should be most important for the child's safety. If a rule is broken, after a short, clear, and gentle explanation, immediately find a place for your child to sit alone for 1 minute. It is very important that punishment come immediately after a rule is broken.

4. Be consistent with discipline. Don't make threats that you cannot carry out. If you say you're going to do it, do it.

Potty training

At 18 months, most toddlers are not yet showing signs that they are ready for toilet training. When toddlers report to parents that they have wet or soiled their diaper, they are beginning to be aware that they prefer dryness. This is a good sign and you should praise your child. Toddlers are naturally curious about the use of the bathroom by other people. Let them watch you or other family members use the toilet. It is important not to put too many demands on a child or shame the child during toilet training.

Safety Tips

- Keep plastic bags, balloons and small hard objects out of reach
- Cut foods into small pieces
- Store toys in a chest without a dropping lid
- Keep hot appliances and cords out of reach
- Don't cook with your child at your feet
- Hold onto your child when you are near traffic
- Provide a play area where balls and riding toys cannot roll into the street
- Always watch your child around any water, including toilets and buckets. Keep toilet seats down and store buckets upside down
- NEVER leave a toddler in a bathtub alone
- Check the stability of drawers, furniture and lamps. Avoid placing furniture (on which children may climb) near windows or on balconies.
- Install window guards on windows and above first floor (unless this is against your local fire codes).
- Make sure windows are closed or have screens that cannot be pushed out. Do not underestimate your child's ability to climb.
- Keep all medicines, vitamins, cleaning fluids locked away.
- Keep the poison control phone number handy: 1-800-222-1222.
- Purchase all medicines with safety caps.

Immunizations

At the 18-month visit, your baby may receive shots. Your baby may run a fever and be irritable for about 1 day after the shots. Your baby may also have some soreness, redness, and swelling in the area where the shots were given. You may give your child acetaminophen drops (1 and 1/2 dropperfuls, or .2 ml, every 4 to 6 hours) to prevent fever.

and irritability. For swelling or soreness, put a wet, v•/arm washcloth on the area of the shots as often and as long as needed for comfort.

Call your child's physician if.

- Your child has a rash or any reaction to the shots other than fever and mild irritability.
- Your child has a fever that lasts more than 36 hours.

Next Visit: Your child's next visit should be at the age of 2 years.